

Check one box and sign below.

I have (for students under the age of 18; My child has):

- had meningococcal immunization within the past five years. The vaccine record is attached.
Note: The Advisory Committee on Immunization Practices recommends that all first-year college students up to age 21 years should have at least one dose of Meningococcal ACWY vaccine not more than 5 years before enrollment, preferably on or after their 16th birthday, and that young adults aged 16 through 23 years may choose to receive the Meningococcal B vaccine series. College students should discuss the Meningococcal B vaccine with a health care provider.
- read or have had explained to me, the information regarding meningococcal disease. I (my child) will obtain immunization against meningococcal disease **within 30 days** from my private health care provider or Farmingdale State College Health and Wellness Center.
- read or have had explained to me, the information regarding meningococcal disease. I understand the risks of not receiving the vaccine. I have decided that I (my child) will **not** obtain immunization against meningococcal disease.

_____ Date _____ / _____ / _____
Signed (Student MUST sign, if under 18, parent or guardian must sign)

_____ Students Date of Birth _____ / _____ / _____
Print Student's Name

_____ Student's E-mail Address _____ Student RAM ID# _____

_____ Student's Mailing Address _____
_____ Student's Phone Number (_____) _____

Return this for to:
**HEALTH & WELLNESS CENTER
FARMINGDALE STATE COLLEGE
2350 BROADHOLLOW ROAD
FARMINGDALE, NY 11735**

About Meningococcal Disease

What is meningococcal disease?

Meningococcal disease is caused by bacteria called Neisseria meningitis. It can lead to serious blood infections. When the linings of the brain and spinal cord become inflamed, it is called meningitis. The disease strikes quickly and can have serious complications, including death. Anyone can get meningococcal disease. Some people are at higher risk. This disease occurs more often in people who are:

- Teenagers or young adults.
- Infants younger than one year of age.
- Living in crowded settings, such as college dormitories or military barracks
- Traveling to areas outside of the United States, such as the "Meningitis belt" in Africa.
- Living with a damaged spleen or no spleen.
- Being treated with Soliris®, or who have complement component deficiency (an inherited immune disorder).
- Exposed during an outbreak.
- Working with meningococcal bacteria in a laboratory.

What are the symptoms?

Symptoms appear suddenly - usually three to four days after a person is infected. It can take up to 10 days to develop symptoms. Symptoms may include:

- A sudden high fever
- Headache
- Stiff neck (meningitis)
- Nausea and vomiting
- Red-purple skin rash
- Weakness and feeling very ill
- Eyes sensitive to light

How is meningococcal disease spread?

It spreads from person-to-person by coughing or coming into close or lengthy contact with someone who is sick or who carries the bacteria. Contact includes kissing, sharing drinks, or living together. Up to one in 10 people carry meningococcal bacteria in their nose or throat without getting sick.

Is there treatment?

Early diagnosis of meningococcal disease is very important. If it is caught early, meningococcal disease can be treated with antibiotics. But, sometimes the infection has caused too much damage for antibiotics to prevent death or serious long-term problems. Most people need to be cared for in a hospital due to serious, life-threatening infections.

What are the complications?

Ten to 15 percent of those who get meningococcal disease die. Among survivors, as many as one in five will have permanent disabilities.

Complications include:

- Hearing loss
- Brain damage
- Kidney damage
- Limb amputations